



Citation/Reference	Mina Mounir and Toon van Waterschoot, Note Onset Detection based on a Spectral Sparsity measure applied to Strings Instruments submitted for DMRN+11: Digital Music Research Network One-day Workshop, London, UK, December 2016.
Archived version	Author manuscript: the content is identical to the content of the published paper, but without the final typesetting by the publisher
Published version	N/A
Journal homepage	http://c4dm.eecs.qmul.ac.uk/dmrn/events/dmrnp11
Author contact	your email mina.mounir@esat.kuleuven.be your phone number + 32 (0) 16-327848
IR	ftp://ftp.esat.kuleuven.be/stadius/mshahata/workshop/NOD_StringInstr_DMRN.pdf

(article begins on next page)



Note Onset Detection based on a Spectral Sparsity measure applied to Strings Instruments

Mina Mounir and Toon van Waterschoot

Department of Electrical Engineering (ESAT), KU Leuven, Belgium,
mina.mounir,toon.vanwaterschoot@esat.kuleuven.be

Abstract— Being the atomic component for a melody’s time dimension, the detection of note onsets is gaining a growing interest in the arising research fields music information retrieval (MIR), machine listening and music processing. We propose a new note onset detection algorithm NINOS² exploiting the spectral sparsity difference between different parts of a musical note. Added to the outstanding performance of NINOS² when applied to automatically annotated guitar melodies and chords progression [1], the proposed algorithm consistently outperforms the state-of-the-art LogSpecFlux (LSF) for the sustained-strings group of instruments crossing the 50% F1-score border. We also propose an additional performance measure to assess the relative position of detected onsets w.r.t. each other.

I. METHOD SUMMARY

In this article, we present NINOS², an algorithm that proposes a new pre-processing and reduction function steps for calculating a smooth Onset Detection Function (ODF) – a highly subsampled version of the original music signal having distinguishable amplitude peaks at time instants where onsets appear [2] – making it easier for the peak picking step to identify the onsets.

For the pre-processing, the subset of low-energy frequency coefficients – traditionally filtered out as they were considered as noise to the detection process – is selected for the ODF calculation. Moreover, NINOS² exploits the difference in spectral sparsity between the transient and the following steady-state component of a musical note when calculating the ODF noted $\aleph(i)$:

$$\aleph(i) = \frac{1}{\sqrt{J}} \frac{\|Y_i\|_2^2}{\|Y_i\|_4^4} \quad (1)$$

where Y is the pre-processed frequency magnitude spectrum, i is the frame index and J is the count of selected coefficients. It is an energy and inverse-sparsity measure as onsets are usually accompanied by an energy increase and

they mark the start of transients which are spectrally less sparse than the steady-state part of a note.

II. EXPERIMENTAL RESULTS

We ran the experiment on a variety of strings instruments: guitar, violin, viola, cello, ...etc. The dataset is generated and automatically annotated as described in [1]. Table I and Fig. 1 show a snippet from the result. In Fig. 1, onsets ground-truth are marked with vertical squares, true positives are marked with circles while false positives with x marks.

TABLE I. PERFORMANCE COMPARISON (BEST-F1 SCORE)

Instruments group	NINOS ²	LSF
Plucked Strings	0.9861	0.9828
Sustained Strings	0.5724	0.4878




Figure 1. Comparison of ODF's and Detections for a Cello.

REFERENCES

- [1] M. Mounir, P. Karsmakers, and T. van Waterschoot, "Guitar note onset detection based on spectral sparsity measure," in *Signal Processing Conference (EUSIPCO), 2016 Proceedings of the 24th European*. IEEE, 2016, pp. 978–982.
- [2] P. Leveau and L. Daudet, "Methodology and tools for the evaluation of automatic onset detection algorithms in music," in *Proc. Int. Symp. Music Information Retrieval*, 2004, pp. 72–75.

This research work was carried out at the ESAT Laboratory of KU Leuven, in the frame of KU Leuven Research Council CoE PFV/10/002 (OPTEC), KU Leuven Impulse Fund IMP/14/037, the FP7-PEOPLE Marie Curie Initial Training Network "Dereverberation and Reverberation of Audio, Music, and Speech (DREAMS)" funded by the European Commission under Grant Agreement no. 316969, IWT O&O Project nr. 150432 "Advances in Auditory Implants: Signal Processing and Clinical Aspects, and IWT O&O Project nr. 150611 "Proof-of-concept of a Rationed Architecture for Vehicle Entertainment and NVH Next-generation Acoustics (RAVENNA). The scientific responsibility is assumed by its authors.